

Product Name: Z-DQMD-FMK Revision Date: 01/10/2021

Product Data Sheet

10

Z-DQMD-FMK

Cat. No.:	A1921	A BOAL
CAS No.:	767287-99-0	top of
Formula:	C29H40FN5O11S	
M.Wt:	685.72	
Synonyms:	Z-DQMD-FMK,Benzyloxycarbonyl-Asp(OMe)-	OF NH O
	GIn-Met-Asp(OMe)-fluoromethylketone	NH ₂
Target:	Apoptosis	049
Pathway:	Caspase	
Storage:	Store at -20°C	010
	DE	OF THE
Colvert 9		Bet

Solvent & Solubility

	\geq 29.2 mg/mL in DMSO; insoluble in EtOH; insoluble in H2O					
In Vitro	Preparing Stock Solutions	Mass Solvent Concentration	1mg	5mg	10mg	
		1 mM	1.4583 mL	7.2916 mL	14.5832 mL	
		5 mM	0.2917 mL	1.4583 mL	2.9166 mL	
		10 mM	0.145 <mark>8 mL</mark>	0.7292 mL	1.4583 mL	

Please refer to the solubility information to select the appropriate solvent.

Biological Activity

IC ₅₀ & Target Cell Viability Assay Cell Line: 3T3-Swiss Albino cells Preparation method: The solubility of this compound in DMSO is > 10 mM. General tips for	C ₅₀ & Target	
Cell Viability Assay Cell Line: 3T3-Swiss Albino cells Preparation method: The solubility of this compound in DMSO is > 10 mM. General tips for	Cell Viability Assay	
Cell Line: 3T3-Swiss Albino cells Preparation method: The solubility of this compound in DMSO is > 10 mM. General tips for	La Applying	
Preparation method: The solubility of this compound in DMSO is > 10 mM. General tips for	Cell Line:	3T3-Swiss Albino cells
a higher concentration: Please warm the tube at 37 °C for 10 minu shake it in the ultrasonic bath for a while. Stock solution can be store 20 °C for several months.	Preparation method:	The solubility of this compound in DMSO is > 10 mM. General tips for obtaining a higher concentration: Please warm the tube at 37 °C for 10 minutes and/or shake it in the ultrasonic bath for a while. Stock solution can be stored below - 20 °C for several months.

1 | www.apexbt.com

	Reacting conditions:	25 μM; 24 hrs	
	Applications:	In zinc-deficient 3T3-Swiss Albino cells, Z-DQMD-FMK (25 μ M) prevented	
		activation of caspase-3. Z-DQMD-FMK treatment did not restore cell number,	
		but resulted in processing of full-length PKC- δ to a 56-kDa fragment.	
In Vivo	Animal experiment		
	Applications:	BIO	

Product Citations

1. Wang J, Shao L, et al. "Synergetic Inhibition of Human Colorectal Cancer Cells by Combining Polyyne-Enriched Fraction from Oplopanax elatus and Irinotecan." Nutr Cancer. 2018 Oct 29:1-11.PMID:30372160

2.Han J, Yang BP, et al. "RhoB/ROCK mediates oxygen-glucose deprivation-stimulatedsyncytiotrophoblast microparticle shedding in preeclampsia." Cell Tissue Res. 2016PMID:27324125

3.Li, Lingfei, et al. "P38/MAPK contributes to endothelial barrier dysfunction via MAP4 phosphorylation-dependent microtubule disassembly in inflammation-induced acute lung injury." Scientific reports 5 (2015).PMID:25746230

See more customer validations on www.apexbt.com.

References

[1]. Susan S. CHOU*, Michael S. CLEGG, Alterations in protein kinase C activity and processing during zinc-deficiency-induced cell death, Biochem. J. (2004) 383, 63–71.

Caution

FOR RESEARCH PURPOSES ONLY.

NOT FOR HUMAN, VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

Specific storage and handling information for each product is indicated on the product datasheet. Most APExBIO products are stable under the recommended conditions. Products are sometimes shipped at a temperature that differs from the recommended storage temperature. Shortterm storage of many products are stable in the short-term at temperatures that differ from that required for long-term storage. We ensure that the product is shipped under conditions that will maintain the quality of the reagents. Upon receipt of the product, follow the storage recommendations on the product data sheet.





PE

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054. Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

2 | www.apexbt.com

