


Product Data Sheet

THZ2

Cat. No.:	A8717
CAS No.:	1604810-84-5
Formula:	C31H28ClN7O2
M.Wt:	566.05
Synonyms:	CDK7-IN-1
Target:	Cell Cycle/Checkpoint
Pathway:	Cyclin-Dependent Kinases
Storage:	Store at -20°C



Solvent & Solubility

≥28.3 mg/mL in DMSO; insoluble in H₂O; ≥2.96 mg/mL in EtOH with gentle warming and ultrasonic

In Vitro	Preparing Stock Solutions	Concentration	Mass		
			Solvent	1mg	5mg
			1 mM	1.7666 mL	8.8331 mL
		5 mM	0.3533 mL	1.7666 mL	3.5333 mL
		10 mM	0.1767 mL	0.8833 mL	1.7666 mL

Please refer to the solubility information to select the appropriate solvent.

Biological Activity

Shortsummary	CDK7 inhibitor
IC ₅₀ & Target	
In Vitro	Cell Viability Assay
	Cell Line: TNBC cells
	Preparation method: The solubility of this compound in DMSO is > 28.3 mg/mL. General tips for obtaining a higher concentration: Please warm the tube at 37 °C for 10 minutes and/or shake it in the ultrasonic bath for a while. Stock solution can be stored below - 20 °C for several months.
Reacting conditions: 0.1 ~ 10000 nM; 7 days	

	Applications:	At low nanomolar doses, THZ2 efficiently suppressed the cell growth of TNBC cells ($IC_{50} = \sim 10$ nM). Besides, THZ2 also induced apoptotic cell death in triple-negative, but not ER/PR+, breast cancer cells or normal human cells. Meanwhile, THZ2 did not elicit an obvious alteration in cell cycle.
Animal experiment		
	Animal models:	Nude mice bearing triple-negative breast tumors
	Dosage form:	10 mg/kg; i.p.; 50 hrs or 25 days
In Vivo	Applications:	In nude mice bearing triple-negative breast tumors, THZ2 significantly reduced the growth rate of tumors without affecting body weight. Moreover, both acute (50 hrs) and chronic exposure (25 days) to THZ2 substantially reduced CTD phosphorylation of RNAPII at all 3 phosphorylation sites (i.e. S2, S5 and S7), which indicated that CDK7 was efficiently inhibited in the tumor cells. According to the immunostaining analysis results of tumor tissues isolated from mice treated with THZ2, reduced proliferation and increased apoptosis were observed.
	Other notes:	Please test the solubility of all compounds indoor, and the actual solubility may slightly differ with the theoretical value. This is caused by an experimental system error and it is normal.

Product Citations

1. Huang JR, Qin WM, et al. "Cyclin-dependent kinase 7 inhibitor THZ2 inhibits the growth of human gastric cancer in vitro and in vivo." Am J Transl Res. 2018 Nov 15;10(11):3664-3676. PMID:30662617

See more customer validations on www.apexbt.com.

References

- [1]. Wang Y, Zhang T, Kwiatkowski N et al. CDK7-dependent transcriptional addiction in triple-negative breast cancer. Cell. 2015 Sep 24;163(1):174-86.

Caution

FOR RESEARCH PURPOSES ONLY.

NOT FOR HUMAN, VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

Specific storage and handling information for each product is indicated on the product datasheet. Most APEXBIO products are stable under the recommended conditions. Products are sometimes shipped at a temperature that differs from the recommended storage temperature. Shortterm storage of many products are stable in the short-term at temperatures that differ from that required for long-term storage. We ensure that the product is shipped under conditions that will maintain the quality of the reagents. Upon receipt of the product, follow the storage recommendations on the product data sheet.

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

