


Product Data Sheet

AEBSF.HCl

Cat. No.:	A2573
CAS No.:	30827-99-7
Formula:	C ₈ H ₁₀ FNO ₂ S·HCl
M.Wt:	239.69
Synonyms:	AEBSF.HCl, AEBSF Hydrochloride
Target:	Serine Protease
Pathway:	Proteases
Storage:	Desiccate at -20°C



Solvent & Solubility

≥12 mg/mL in DMSO, ≥15.73 mg/mL in H₂O, ≥23.8 mg/mL in EtOH with gentle warming

In Vitro

	Solvent Concentration	Mass	1mg	5mg	10mg
Preparing Stock Solutions	1 mM		4.1721 mL	20.8603 mL	41.7206 mL
	5 mM		0.8344 mL	4.1721 mL	8.3441 mL
	10 mM		0.4172 mL	2.0860 mL	4.1721 mL

Please refer to the solubility information to select the appropriate solvent.

Biological Activity

Shortsummary

Serine protease inhibitor

IC₅₀ & Target

In Vitro

Cell Viability Assay

Cell Line:	K695sw, HS695 and SKN695 cells
Preparation method:	The solubility of this compound in DMSO is ≥798.97mg/mL. General tips for obtaining a higher concentration: Please warm the tube at 37°C for 10 minutes and/or shake it in the ultrasonic bath for a while. Stock solution can be stored below -20°C for several months.
Reacting conditions:	0, 0.1, 0.3, 0.6, 0.8, 1.0 and 1.2 mM; 20 mins
Applications:	AEBSF was found to inhibit Aβ production in various cell lines. In K293 cells

transfected with β APP695 (K695sw), AEBSF showed dose-dependent reduction of A β with the IC50 value of about 1 mM. In HS695 and SKN695 cells transfected with wild-type APP695, AEBSF showed inhibition effect with IC50 value of about 300 μ M. AEBSF was also found to increase α -cleavage and inhibit β - cleavage.

Animal experiment

Animal models:	SD rats
Dosage form:	Intrauterine (5 mg or 10 mg AEBSF per injection) or tail vein (10 mg AEBSF per rat) administration on day 3 of pregnancy
Applications:	According to the number of visible implanted embryos on day 8 of pregnancy, it was shown that AEBSF inhibited embryo implantation in rat.
Other notes:	Please test the solubility of all compounds indoor, and the actual solubility may slightly differ with the theoretical value. This is caused by an experimental system error and it is normal.

In Vivo

Product Citations

1. Brodie NI, Popov KI, et al."Conformational ensemble of native α -synuclein in solution as determined by short-distance crosslinking constraint-guided discrete molecular dynamics simulations." PLoS Comput Biol. 2019 Mar 27;15(3):e1006859.PMID:30917118

See more customer validations on www.apexbt.com.

References

- [1]. Citron M, Diehl T S, Capell A, et al. Inhibition of amyloid β -protein production in neural cells by the serine protease inhibitor AEBSF. Neuron, 1996, 17(1): 171-179.
- [2]. Jiang YH, Shi Y, He YP, Du J, Li RS, Shi HJ, Sun ZG, Wang J. Serine protease inhibitor 4-(2-aminoethyl)benzenesulfonyl fluoride hydrochloride (AEBSF) inhibits the rat embryo implantation in vivo and interferes with cell adhesion in vitro. Contraception. 2011 Dec;84(6):642-8.

Caution

FOR RESEARCH PURPOSES ONLY.

NOT FOR HUMAN, VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

Specific storage and handling information for each product is indicated on the product datasheet. Most APExBIO products are stable under the recommended conditions. Products are sometimes shipped at a temperature that differs from the recommended storage temperature. Shortterm storage of many products are stable in the short-term at temperatures that differ from that required for long-term storage. We ensure that the product is shipped under conditions that will maintain the quality of the reagents. Upon receipt of the product, follow the storage recommendations on the product data sheet.

APEX BIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

