

FITC Goat Anti-Rabbit IgG (H+L) Antibody

General Information

| | |
|---------------------|---|
| Name | FITC Goat Anti-Rabbit IgG (H+L) Antibody |
| Host species | Goat |
| Target species | Rabbit |
| Tested applications | Suitable for: ICC/IF, IHC-Fr, IHC-P, ICC, Flow Cyt, ELISA, WB |
| Conjugation | FITC. Ex: 495 nm, Em: 519 nm |


Properties

| | |
|----------------------|---|
| Form | Liquid |
| Storage buffer | Preservative: 0.02% Sodium azide Constituents: 23% Glycerol, PBS, 1% BSA |
| Storage instructions | Shipped at 4° C. Store at 4° C short term (2 weeks). Upon delivery aliquot. Store at -20° C. Avoid freeze / thaw cycle. Stable for 12 months at -20° C. Keep away from Light. |
| Concentration | 1 mg/mL |
| Purity | Immunogen affinity purified |
| Purification notes | This antibody was isolated by affinity chromatography using antigen coupled to agarose beads. |
| Clonality | Polyclonal |
| Isotype | IgG |

Applications

| Application | Notes |
|-------------|-------------------------------------|
| ICC/IF | 1/500 - 1/2000 |
| IHC-Fr | Use at an assay dependent dilution. |
| IHC-P | 1/100-1/500 |
| Flow Cyt | 1/250 - 1/1000 |
| ELISA | Use at an assay dependent dilution. |
| WB | 1/20-1/50 |

Images



Immunofluorescence analysis of HeLa cells, using PHB antibody as the primary antibody. The cells were incubated with the primary antibody overnight at 4° C.

Secondary antibody: FITC Goat Anti-Rabbit IgG (H+L) Antibody at dilution of 1:1000.

Blue: DAPI for nuclear staining.

Notes

Our product is suitable for single-labeling experiment, to conduct multi-labeling experiments, secondary antibodies with pre-affinity treatment of serum proteins or immunoglobulin of close species are recommended. .

APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com