

Western Stripping Buffer (Neutral)

Introduction

Western Blot Stripping Buffer is used for the reuse of membranes onto which proteins have been transferred during Western blotting. After completing primary and secondary antibody binding and subsequent chemiluminescent detection, it is sometimes necessary to detect reference proteins with relatively stable expression levels, such as tubulin or actin, or to detect other proteins for comparison. By using this stripping buffer to thoroughly remove the primary and secondary antibodies, the used membrane can be conveniently reused to detect other proteins. Compared to running a new SDS-PAGE gel, this method is not only time-saving and labor-saving but also eliminates errors caused by reloading samples, thereby improving comparability.

Reusing the same membrane multiple times with Western Blot Stripping Buffer may lead to a reduction in protein signal. However, this product has been tested with various antibodies and typically allows for 3–5 reuses of the membrane.

This reagent takes only approximately 20–30 minutes to enable membrane reuse, after which you can proceed with blocking and subsequent Western blotting steps.


Different Western blot stripping buffers vary slightly in their principles for removing bound primary and secondary antibodies. This product utilizes different combinations of detergents, acidic agents, alkaline agents, reducing agents, chelators, salt concentrations, and other specialized reagents to disrupt primary and secondary antibody binding without causing significant loss of the proteins transferred onto the membrane.

Protocol

1. After completing Western chemiluminescent detection, rinse the membrane in distilled water for 5 minutes.
2. Discard the distilled water and add an appropriate amount of Western Stripping Buffer (Neutral), ensuring the membrane is fully covered. Shake on an orbital shaker for 30–60 minutes.
3. Discard the Western Stripping Buffer (Neutral) and remove any residual liquid. Rinse the membrane 3–4 times with TBS, TBST, or PBS, shaking for 3–5 minutes each time on an orbital shaker.
4. Proceed with blocking and subsequent Western blot steps.

■ Note

1. When using horseradish peroxidase (HRP), any blocking step should be performed with 5% non-fat milk or BSA. If using alkaline phosphatase, any blocking step should be performed with casein.
2. This Western Blot Stripping Buffer is suitable for PVDF membranes only. Using nitrocellulose membranes will result in significant loss of transferred proteins.
3. This reagent is suitable for Western detection using ECL chemiluminescent reagents. It is not suitable for Western detection using non-chemiluminescent methods such as DAB or NBT/BCIP.
4. Storage conditions: Store at 4°C; stable for two months. Store at -20°C; stable for one year.
5. This product is for scientific use only.



APEX BIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

