

Anti-Acetyl-Histone H4 (Lys5) Rabbit Monoclonal Antibody

Introduction

The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family.

Product parameters

Alternative Names	H4K5ac; H4F2; HIST; HIST1H4A; Histone H4.AC-H4K5; H4K5
Gene ID	121504
Gene Name	H4C1
SwissProt ID	P62805
Host	Rabbit
Reactivity	Human, Mouse
Molecular Weight	Calculated MW: 11 kDa; Observed MW: 11 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Acetylated
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-5D6F5
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthesized peptide derived from human Histone H4 (acetyl K5)
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB, IHC-F, IHC-P, ICC/IF, IP
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/20
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Shipping	Blue ice

Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

