

Anti-ACSS2 Rabbit Monoclonal Antibody

Activates acetate so that it can be used for lipid synthesis or for energy generation. Cytoplasmic acetyl-CoA synthetase (AceCS1) catalyzes the conversion of acetate and CoA to acetyl-CoA. Acetyl-CoA synthesized by AceCS1 is used for fatty acid and lipid biosynthesis. Studies suggest that this enzyme is regulated by sterol regulatory element-binding proteins.

Product parameters

Introduction

Alternative Names	ACSS2; ACSA; ACAS2; AceCS; MYH7B; Acetyl CoA synthetase
Gene ID	55902
Gene Name	ACSS2
SwissProt ID	Q9NR19
Host	Rabbit
Reactivity	
Molecular Weight	Calculated MW: 79 kDa; Observed MW: 79 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-4C12G3
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human ACSS2
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB
Dilution Ratio	WB: 1/500-1/1000
Research Field	Signal Transduction
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol P Ex B 0



Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





