

Anti-Activin Receptor Type IA Rabbit Monoclonal Antibody

Introduction

On ligand binding, forms a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators.

Product parameters

| Alternative Names | ACTRI; Acvr1; ACVR1A; ACVRLK2; ALK2; FOP; SKR1; TSRI |
|--------------------------------|---|
| Gene ID | 90 |
| Gene Name | ACVR1 |
| SwissProt ID | Q04771 |
| Host | Rabbit |
| Reactivity | Human, Mouse |
| Mole <mark>cular</mark> Weight | Calculated MW: 57 kDa; Observed MW: 57 kDa |
| Conjugation | Unconjugated Achieve Perfection, Explore the Unknown |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-20B12E2 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Not Free |
| Immunogen | A synthetic peptide of human Activin Receptor Type IA |
| Purification | Affinity Purified |
| Buffer System | 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA. |
| Application | WB, IP |
| Dilution Ratio | WB: 1/500-1/1000 IP: 1/20 |
| Research Field | Signal Transduction |
| Product Categories | Primary antibody |
| Shipping | Blue ice |

| Storage | -20°C |
|-----------------|----------------------------------|
| Expiration Date | 12 months |
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





