

## Anti-Angiopoietin 2 Rabbit Monoclonal Antibody

### Introduction

Can induce tyrosine phosphorylation of TIE2. Binds to TIE2 receptor and counteracts blood vessel maturation/stability mediated by angiopoietin-1. Its function may be context-dependent.

### Product parameters

Alternative Names	Angiopoietin-2 (ANG-2)
Gene ID	285
Gene Name	ANGPT2
SwissProt ID	O15123
Host	Rabbit
Reactivity	Rat
Molecular Weight	Calculated MW: 57 kDa; Observed MW: 57 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-19A8C5
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human Angiopoietin 2
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB, IHC-P
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100
Research Field	Cardiovascular
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C

Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

## Protocol

Configure the product according to the application range and recommended dilution ratio.

**\*Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.



**APExBIO Technology**

**[www.apexbt.com](http://www.apexbt.com)**

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: [info@apexbt.com](mailto:info@apexbt.com)

