

Anti-APAF1 Rabbit Monoclonal Antibody

Introduction

APAF Oligomeric Apaf-1 mediates the cytochrome c-dependent autocatalytic activation of pro-caspase-9 (Apaf-3), leading to the activation of caspase-3 and apoptosis. This activation requires ATP. Isoform 6 is less effective in inducing apoptosis. Induced by E2F and p53 in apoptotic neurons. Monomer. Oligomerizes upon binding of cytochrome c and dATP. Oligomeric Apaf-1 and pro-caspase-9 bind to each other via their respective NH2-terminal CARD domains and consecutively mature caspase-9 is released from the complex. Pro-caspase-3 is recruited into the Apaf-1-pro-caspase-9 complex via interaction with pro-caspase-9. Interacts with APIP. 6 isoforms of the human protein are produced by alternative splicing.

Product parameters

| | |
|-------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Alternative Names | APAF; APAF-1; APAF1; apoptotic peptidase activating factor 1; apoptotic protease activating factor 1; Apoptotic protease-activating factor 1; CED4; DKFZp781B1145; KIAA0413 |
| Gene ID | 317 |
| Gene Name | APAF1 |
| SwissProt ID | O14727 |
| Host | Rabbit |
| Reactivity | Human, Mouse |
| Molecular Weight | Calculated MW: 142 kDa; Observed MW: 135 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-15A1H3 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Free |
| Immunogen | A synthesized peptide derived from human APAF1 |
| Purification | Affinity Chromatography |
| Buffer System | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Application | WB, IHC-P, ICC/IF |

| | |
|--------------------|-------------------------------------------------|
| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 |
| Research Field | Cell Biology |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |
| Expiration Date | 12 months |
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

