

Anti-Aspartate Aminotransferase Rabbit Monoclonal Antibody

Introduction

Plays a key role in amino acid metabolism. Important for metabolite exchange between mitochondria and cytosol. Facilitates cellular uptake of long-chain free fatty acids.

Product parameters

| Alternative Names | KAT4; KATIV; KYAT4; mitAAT |
|----------------------------|---|
| Gene ID | 2806 |
| Gene Name | GOT2 |
| SwissProt ID | P00505 |
| Host | Rabbit |
| Reactivity | Human, Mouse, Rat |
| Molecular Weight | Calculated MW: 48 kDa; Observed MW: 45 kDa |
| Conjugation | Unconjugated |
| Ex Achieve Perfect | Achieve Perfection, Explore the Unknown |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-8G8D2 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Not Free |
| Immunogen | A synthetic peptide of human FABP-1 |
| Purification | Affinity Purified |
| Buffer System | 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA. |
| Ap <mark>plica</mark> tion | WB, IHC-P, IP |
| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20 |
| Research Field | Signal Transduction |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |
| | |

| Expiration Date | 12 months |
|-----------------|----------------------------------|
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





