Anti-beta 2 Microglobulin (4C3) Mouse Monoclonal Antibody

Introduction

Major histocompatibility complex (MHC) class 1 molecules bind to antigens for presentation on the surface of cells. The proteasome is responsible for producing these antigens from the components of foreign pathogens. MHC class 1 molecules consist of an a heavy chain that contains three subdomains (α1, α2, α3), and a non-covalent associating light chain, known as β-2-Microglobulin.

Product parameters

Alternative Names	B2MG; Beta 2 microglobin; Beta 2 microglobulin; Beta-2-microglobulin form pI 5.3; CDABP0092; Hdcma22p
Gene ID	567
Gene Name	B2M
SwissProt ID	P61769
Host	Mouse
Reactivity	Human
Molecular Weight	-
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG2b
Isotype	Monoclonal Antibody
Clonality No.	AP-17H2A12
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Purified recombinant human β2-microglobulin protein expressed in E.coli.
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	ELISA
Dilution Ratio	ELISA: 1/10000
Research Field	Cardiovascular
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.

APExBIO Technology
www.apexbt.com
7505 Fannin street, Suite 410, Houston, TX 77054.
Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com