

Anti-beta 2 Microglobulin Rabbit Monoclonal Antibody

Introduction

Major histocompatibility complex (MHC) class 1 molecules bind to antigens for presentation on the surface of cells. The proteasome is responsible for producing these antigens from the components of foreign pathogens. MHC class 1 molecules consist of an a heavy chain that contains three subdomains (α 1, α 2, α 3), and a non-covalent associating light chain, known as β -2-Microglobulin.

Product parameters

Alternative Names	B2MG; Beta 2 microglobin; Beta 2 microglobulin; Beta-2-microglobulin form pl 5.3; CDABP0092; Hdcm
Gene ID	567
Gene Name	B2M
SwissProt ID	P61769
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 14 kDa; Observed MW: 14 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-3A6F3
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human beta 2 Microglobulin
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, ICC/IF, IP
Dilution Ratio	WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/30
Research Field	Cancer
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

| Protocol | | |



Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





