

Anti-CD105 Rabbit Monoclonal Antibody

Introduction

ainding of andatholial calls to integrins

Major glycoprotein of vascular endothelium. May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors.

Product parameters

| Alternative Names | ENG; END; Endoglin; CD antigen CD105 |
|---------------------|---|
| Gene ID | 2022 |
| Gene Name | ENG |
| SwissProt ID | P17813 |
| Host | Rabbit |
| Reactivity | Human |
| Molecular Weight | Calculated MW: 71 kDa; Observed MW: 71 kDa |
| Conjugation | Unconjugated PEYB |
| Ex Achieve Perfecti | Achieve Perfection, Explore the Unknown |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-2F12H5 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Free |
| Immunogen | A synthesized peptide derived from human CD105 |
| Purification | Affinity Chromatography |
| Buffer System | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Application | WB, IHC-P |
| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 |
| Research Field | Immunology |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |
| | |

| Expiration Date | 12 months |
|-----------------|----------------------------------|
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





