

Anti-CD81 Rabbit Monoclonal Antibody

Introduction

The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. Many research studies demonstrate a role for CD81 in lymphocyte signaling. CD81 is also a well-characterized receptor for Hepatitis C Virus and facilitates the entry of the virus into target cells.

Product parameters

Alternative Names	CD81; TAPA1; TSPAN28; CD81 antigen; 26 kDa cell surface protein TAPA-1; Target of the antiproliferative antibody 1; Tetraspanin-28; Tspan-28; CD81
Gene ID	975
Gene Name	CD81
SwissProt ID	P60033
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 26 kDa; Observed MW: 26 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-17F7F11
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human CD81
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, ICC/IF, FC
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 FC: 1/50-1/100
Research Field	Immunology
Product Categories	Primary antibody
Shipping	Blue ice

Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

