

Anti-CHD3 (7F6) Mouse Monoclonal Antibody

Introduction

Component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin by deacetylating histones. Required for anchoring centrosomal pericentrin in both interphase and mitosis, for spindle organization and centrosome integrity.

Product parameters

Alternative Names	CHD3; Chromodomain-helicase-DNA-binding protein 3; CHD-3; ATP-dependent helicase CHD3; Mi-2 autoantigen 240 kDa protein; Mi2-alpha; Zinc finger helicase; hZFH
Gene ID	1107
Gene Name	CHD3
SwissProt ID	Q12873
Host	Mouse
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 227 kDa; Observed MW: 260 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG1
Isotype	Monoclonal Antibody
Clonality No.	AP-10F3B12
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Purified recombinant human CHD3 (C-terminus) protein fragments expressed in E.coli.
Purification	Affinity purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	WB, ICC/IF, IP
Dilution Ratio	WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/20
Research Field	Immunology
Product Categories	Primary antibody
Shipping	Blue ice

Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APEx BIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com