

Anti-Cleaved-Caspase 2 Rabbit Monoclonal Antibody

Introduction

Caspase-2 (Nedd2, ICH-1) is an aspartate-specific cysteine protease that is activated in response to various apoptotic stimuli. Caspase-2 is unique among the caspases in that it has features of both upstream caspases (long prodomain) and downstream caspases (DEXD substrate specificity). Caspase-2 is highly expressed in the brain during development and is expressed at low levels in adult tissue.

Product parameters

Alternative Names	Caspase-2; Caspase 2; Caspase2; CASP-2; Neural precursor cell expressed developmental down-regulated protein 2; NEDD-2; Protease ICH-1; CASP2; ICH1; NEDD2
Gene ID	835
Gene Name	CASP2
SwissProt ID	P42575
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 51 kDa; Observed MW: 48,30,12 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-5H9D6
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Im <mark>mun</mark> ogen	A synthesized peptide derived from human Caspase-2
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, ICC/IF, FC
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 FC: 1/50-1/100
Research Field	Cell Biology
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

■ Protocol → B



Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





