

## **Anti-Cofilin Rabbit Monoclonal Antibody**

## Introduction



The protein encoded by this gene can polymerize and depolymerize F-actin and G-actin in a pH-dependent manner. Increased phosphorylation of this protein by LIM kinase aids in Rho-induced reorganization of the actin cytoskeleton. Cofilin is a widely distributed intracellular actin-modulating protein that binds and depolymerizes filamentous F-actin and inhibits the polymerization of monomeric G-actin in a pH-dependent manner. It is involved in the translocation of actin-cofilin complex from cytoplasm to nucleus.[supplied by OMIM, Apr 2004]

## Product parameters

Alternative Names	CFL1; CFL; Cofilin-1; 18 kDa phosphoprotein; p18; Cofilin; non-muscle isoform	
Gene ID	1072	
Gene Name	CFL1	
SwissProt ID	P23528	
Host	Rabbit	
Reactivity	Human	
Molecular Weight	Calculated MW: 19 kDa; Observed MW: 19 kDa	
Conjugation	Unconjugated	
Ex	-	
Em	-	
Modification	Unmodified	
Clonality	IgG	
Isotype	Monoclonal Antibody	
Clonality No.	AP-19E9B6	
Form	Liquid	
Concentration	See label	
Carrier	Carrier Not Free	
Immunogen	A synthetic peptide of human Cofilin	
Purification	Affinity Purified	
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.	
Application	WB, IHC-F, IHC-P, ICC/IF	
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200	
Research Field	Signal Transduction	

Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.





