


Anti-Cullin 5 Rabbit Monoclonal Antibody

Introduction



proteosomal degradation of p53/TP53 stimulated by adenovirus E1B-55 kDa protein. May form a cell surface

Product parameters

vasopressin receptor.

Gene Name CUL5 SwissProt ID Q93034 Host Rabbit Reactivity Human, Mouse, Rat Molecular Weight Calculated MW: 91 kDa; Observed MW: 91 kDa Conjugation Unconjugated Ex - Em - Modification Unmodified Clonality IgG Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
SwissProt ID Q93034 Host Rabbit Reactivity Human, Mouse, Rat Molecular Weight Calculated MW: 91 kDa; Observed MW: 91 kDa Conjugation Unconjugated Ex - Em - Modification Unmodified Clonality IgG Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
Host Rabbit Reactivity Human, Mouse, Rat Molecular Weight Calculated MW: 91 kDa; Observed MW: 91 kDa Conjugation Unconjugated Ex - Em - Modification Unmodified Clonality IgG Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
Reactivity Human, Mouse, Rat Molecular Weight Calculated MW: 91 kDa; Observed MW: 91 kDa Conjugation Unconjugated Ex - Em - Modification Unmodified Clonality IgG Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
Molecular Weight Calculated MW: 91 kDa; Observed MW: 91 kDa Conjugation Unconjugated Ex - Em - Modification Unmodified Clonality IgG Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
Conjugation Unconjugated Ex - Em - Modification Unmodified Clonality IgG Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
Ex - Em - Modification Unmodified Clonality IgG Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
Em - Modification Unmodified Clonality IgG Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
Modification Unmodified Clonality IgG Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
Clonality IgG Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
Isotype Monoclonal Antibody Clonality No. AP-3C11F10 Form Liquid	
Clonality No. AP-3C11F10 Form Liquid	
Form Liquid	
A DELDIA	
Concentration See label	
Carrier Not Free	
Immunogen Recombinant protein of human Cullin 5	
Purification Affinity Purified	
Buffer System 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.	
Application WB	

Dilution Ratio	WB: 1/500-1/1000
Research Field	Cell Biology
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.

















