

Anti-Cyclophilin B Rabbit Monoclonal Antibody

Introduction

PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides.

Product parameters

Alternative Names	PPIB; CYPB; Peptidyl-prolyl cis-trans isomerase B; PPlase B; CYP-S1; Cyclophilin B; Rotamase B
	S-cyclophilin; SCYLP
Gene ID	5479
Gene Name	PPIB
SwissProt ID	P23284
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 24 kDa; Observed MW: 20 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-11G11C6
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human Cyclophilin B
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB In Unknown
Dilution Ratio	WB: 1/500-1/1000
Research Field	Immunology
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
J	

Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





