

## Anti-Cystatin C (3B11) Mouse Monoclonal Antibody

### Introduction

Cystatin C is a 14 kDa member of the Cystatin superfamily of cysteine protease inhibitors. Most cell types secrete Cystatin C. Cystatin C inhibits cathepsins, and thereby may function as a tumor suppressor by inhibiting cathepsin mediated tumor cell invasion. In addition, this tumor suppressor function can also be attributed to Cystatin C's ability to antagonize TGF- $\beta$ 1 signaling.

### Product parameters

Alternative Names	CST3; Cystatin-C; Cystatin-3; Gamma-trace; Neuroendocrine basic polypeptide; Post-gamma-globulin
Gene ID	1471
Gene Name	CST3
SwissProt ID	P01034
Host	Mouse
Reactivity	Human
Molecular Weight	-
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG1
Isotype	Monoclonal Antibody
Clonality No.	AP-8D2F9
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Recombinant human cystatin c protein.
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	ELISA
Dilution Ratio	ELISA: 1/10000
Research Field	Tags & Cell Markers
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

## Protocol



Configure the product according to the application range and recommended dilution ratio.

**\*Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.



**APExBIO Technology**

**[www.apexbt.com](http://www.apexbt.com)**

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: [info@apexbt.com](mailto:info@apexbt.com)

