

Anti-Dopamine Transporter Rabbit Monoclonal Antibody

Introduction

Amine transporter. Terminates the action of dopamine by its high affinity sodium-dependent reuptake into presynaptic terminals.

Product parameters

| Alternative Names | DA transporter; DAT1; Dopamine transporter 1; PKDYS; SLC6A3 |
|--------------------|---|
| Gene ID | 6531 |
| Gene Name | SLC6A3 |
| SwissProt ID | Q01959 |
| Host | Rabbit |
| Reactivity | Human |
| Molecular Weight | Calculated MW: 68 kDa; Observed MW: 50 kDa |
| Conjugation | Unconjugated A P E B |
| Ex Achieve Perfect | en, Explore the Unknown. Achieve Perfection, Explore the Unknown. |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-16A7A3 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Free |
| Immunogen | A synthesized peptide derived from human Dopamine Transporter |
| Purification | Affinity Chromatography |
| Buffer System | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Application | WB, FC, IP |
| Dilution Ratio | WB: 1/500-1/1000 IP: 1/50 FC: 1/50-1/100 |
| Research Field | Neuroscience |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |
| | |

| Expiration Date | 12 months |
|-----------------|----------------------------------|
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





