

Anti-EHMT2 Rabbit Monoclonal Antibody

Introduction

Histone methyltransferase that specifically mono- and dimethylates 'Lys-9' of histone H3 (H3K9me1 and H3K9me2, respectively) in euchromatin. H3K9me represents a specific tag for epigenetic transcriptional repression by recruiting HP1 proteins to methylated histones.

Product parameters

| | |
|--------------------|---|
| Alternative Names | Bat8; Ehmt2; G9A; GAT8; NG36 |
| Gene ID | 10919 |
| Gene Name | EHMT2 |
| SwissProt ID | Q96KQ7 |
| Host | Rabbit |
| Reactivity | Human |
| Molecular Weight | Calculated MW: 132 kDa; Observed MW: 170 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-1E5A1 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Free |
| Immunogen | A synthesized peptide derived from human EHMT2/G9A |
| Purification | Affinity Purified |
| Buffer System | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Application | WB, IHC-P |
| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 |
| Research Field | Epigenetics and Nuclear Signaling |
| Product Categories | Primary antibody |
| Shipping | Blue ice |

| | |
|-----------------|----------------------------------|
| Storage | -20°C |
| Expiration Date | 12 months |
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

