

## **Anti-eIF4E Rabbit Monoclonal Antibody**

## Introduction

eIF4E, a protein modulates translation of maternal mRNAs in early embryos before the onset of zygotic transcription. eIF4E also influences the overall rate of translation. eIF4E binds to the 7 methyl GTP cap structure of eukaryotic mRNAs. Phosphorylation of eIF4E on serine 209 regulates the affinity of this protein for the 7 methyl GTP cap and/or RNA. Phosphorylation also enhances the interaction of eIF4E with eIF4G, which form a complex known as eIF4F. eIF4E phosphorylation is correlated with increased translational rate in a number of cell types.

## Product parameters

Alternative Names	EIF4E; EIF4EL1; EIF4F; Eukaryotic translation initiation factor 4E; eIF-4E; eIF-4F 25 kDa subul mRNA cap-binding protein
Gene ID	1977
Gene Name	EIF4E
SwissProt ID	P06730
Host Achieve Perfect	Rabbit
Reactivity	Human
Molecular Weight	-
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-19C2E12
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen Period	A synthesized peptide derived from human elF4E
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	IHC-P
Dilution Ratio	IHC: 1/100-1/200
Research Field	Epigenetics and Nuclear Signaling

Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.





