

Anti-EMAP II Rabbit Monoclonal Antibody

Introduction

Non-catalytic component of the multisynthase complex. Stimulates the catalytic activity of cytoplasmic arginyl-tRNA synthase. Binds tRNA. Possesses inflammatory cytokine activity. Negatively regulates TGF-beta signaling through stabilization of SMURF2 by binding to SMURF2 and inhibiting its SMAD7-mediated degradation. Involved in glucose homeostasis through induction of glucagon secretion at low glucose levels. Promotes dermal fibroblast proliferation and wound repair. Regulates KDELR1-mediated retention of HSP90B1/gp96 in the endoplasmic reticulum. Plays a role in angiogenesis by inducing endothelial cell migration at low concentrations and endothelian cell apoptosis at high concentrations. Induces maturation of dendritic cells and monocyte cell adhesion. Modulates endothelial cell responses by degrading HIF-1A through interaction with PSMA7.

Product parameters

| Alternative Names | p43; HLD3; EMAP2; SCYE1; EMAPII; AIMP1 |
|-------------------|---|
| Gene ID | 9255 O |
| Gene Name | AIMP1 Achieve Perfection, Explore the Unknown |
| SwissProt ID | Q12904 |
| Host | Rabbit |
| Reactivity | Human, Rat |
| Molecular Weight | Calculated MW: 34 kDa; Observed MW: 34 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-10F4E7 |
| Form | Liquid O APEX 5 O |
| Concentration | See label |
| Carrier | Carrier Not Free |
| Immunogen | A synthetic peptide of human EMAP II |
| Purification | Affinity Purified |
| Buffer System | 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA. |

| Application | WB |
|-------------------------------|----------------------------------|
| Dilution Ratio | WB: 1/500-1/1000 |
| Research Field | Cell Biology |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |
| Expi <mark>ration</mark> Date | 12 months APEXBO |
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.

















