


Anti-FES Rabbit Monoclonal Antibody

Introduction



actin cytoskeleton, microtubule assembly, cell attachment and cell spreading. Plays a role in FCER1 (high affinity immunoglobulin epsilon receptor)-mediated signaling in mast cells. Acts down-stream of the activated FCER1 receptor and the mast/stem cell growth factor receptor KIT. Plays a role in the regulation of mast cell degranulation. Plays a role in the regulation of cell differentiation and promotes neurite outgrowth in response to NGF signaling. Plays a role in cell scattering and cell migration in response to HGF-induced activation of EZR. Phosphorylates BCR and down-regulates BCR kinase activity. Phosphorylates HCLS1/HS1, PECAM1, STAT3 and TRIM28.

Product parameters

Alternative Names	FPS
Gene ID	2242 O
Gene Name	FES Achieve Perfection, Explore the Unknown
SwissProt ID	P07332
Host	Rabbit
Reactivity	Human
Molecular Weight	Calculated MW: 93 kDa; Observed MW: 93 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-12A1H8
Form	Liquid O APEXBIO
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human FES
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.

Application	WB	
Dilution Ratio	WB: 1/500-1/1000	
Research Field	Signal Transduction	
Product Categories	Primary antibody	
Shipping	Blue ice	
Storage	-20°C	
Expi <mark>ration</mark> Date	12 months APEXBO	
Note	Please avoid freeze-thaw cycles.	

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.

















