


Anti-FRA2 Rabbit Monoclonal Antibody

Introduction



Controls osteoclast survival and size (By similarity). As a dimer with JUN, activates LIF transcription (By similarity). Activates CEBPB transcription in PGE2-activated osteoblasts.

Product parameters

Alternative Names	FOSL2; FRA2; Fos-related antigen 2; FRA-2
Gene ID	2355
Gene Name	FOSL2
SwissProt ID	P15408
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 35 kDa; Observed MW: 45 kDa
Conjugation	Unconjugated PEXBIO
Ex Achieve Perfect	Epitore the Unknown.
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-14G12G10
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human FRA2
Purification	Affinity Purified
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, FC, IP
Dilution Ratio	WB: 1/500-1/1000 IP: 1/50 FC: 1/50-1/100
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C

Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





