

## Anti-Frizzled 9 Rabbit Monoclonal Antibody

### Introduction

Receptor for Wnt proteins. Most of frizzled receptors are coupled to the beta-catenin canonical signaling pathway, which leads to the activation of disheveled proteins, inhibition of GSK-3 kinase, nuclear accumulation of beta-catenin and activation of Wnt target genes.

### Product parameters

Alternative Names	CD349; frizzled-9; Fz-9; FZD3; FzE6
Gene ID	8326
Gene Name	FZD9
SwissProt ID	O00144
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 64 kDa; Observed MW: 64 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-1D8D1
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human Frizzled 9 / CD349
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB
Dilution Ratio	WB: 1/500-1/1000
Research Field	Neuroscience
Product Categories	Primary antibody
Shipping	Blue ice

Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

## Protocol

Configure the product according to the application range and recommended dilution ratio.

**\*Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.



**APExBIO Technology**

**[www.apexbt.com](http://www.apexbt.com)**

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: [info@apexbt.com](mailto:info@apexbt.com)

