

Anti-HDAC11 Rabbit Monoclonal Antibody

Introduction

play an important role in the modification of chromatin structure and thus in the suppression and activation of transcription and cellular differentiation. There are 11 members in the HDAC family that are classified into four classes. Class I HDACs represent homologs of the yeast histone deacetylase RPD3, class II HDACs share strong homology with the yeast histone deacetylase HDA1, class III HDAC are closely related to the yeast SIR2 protein, and class IV HDACs comprises Histone deacetylase 11 (HDAC11)-related enzymes. HDAC11 contains 347 amino acid residues.

Product parameters

Alternative Names	HDAC11; HD11; Histone deacetylase 11
Gene ID	79885
Gene Name	HDAC11
SwissProt ID	Q96DB2
Host	Rabbit
Reactivity	Human
Molecular Weight	Calculated MW: 39 kDa; Observed MW: 39 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-14B10D10
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human HDAC11
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IP
Dilution Ratio	WB: 1/500-1/1000 IP: 1/50

Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.



Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

