

Anti-HDAC3 (3G3) Mouse Monoclonal Antibody

Introduction

HDAC3 is a nuclear and cytoplasmic protein that deacetylates both histone (H2A, H3, H4) and non-histone substrates (RelA, SRY, p53, MEF2, PCAF and p300/CBP). HDAC3 deacetylase activity is stimulated by interactions with the N-CoR and SMRT co-repressor proteins. Together, these three proteins form a functional complex that represses transcription associated with nuclear hormone receptors and other transcription factors, including Rev-Erb, COUP-TF, DAX1, MAD and Pit-1.

Product parameters

Alternative Names	HDAC3; Histone deacetylase 3; HD3; RPD3-2; SMAP45
Gene ID	8841
Gene Name	HDAC3
SwissProt ID	O15379
Host	Mouse
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 49 kDa; Observed MW: 49 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG1
Isotype	Monoclonal Antibody
Clonality No.	AP-8F2F1
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Purified recombinant human HDAC3 protein fragments expressed in E.coli.
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	WB
Dilution Ratio	WB: 1/500-1/1000
Research Field	Epigenetics and Nuclear Signaling

Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.



Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

