

Anti-Histone H1.2 Rabbit Monoclonal Antibody

Introduction

Histones H1 are necessary for the condensation of nucleosome chains into higher order structures.

Product parameters

| | |
|--------------------|---|
| Alternative Names | H1.a; H1F2; H1s-1; HIST1H1C; Histone H1.2; Histone H1c; Histone H1d; Histone H1s-1 |
| Gene ID | 3006 |
| Gene Name | H1-2 |
| SwissProt ID | P16403 |
| Host | Rabbit |
| Reactivity | Human, Mouse, Rat |
| Molecular Weight | Calculated MW: 21 kDa; Observed MW: 29 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-3A3D8 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Free |
| Immunogen | A synthesized peptide derived from human Histone H1.2 |
| Purification | Affinity Chromatography |
| Buffer System | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Application | WB, IHC-P, ICC/IF, ChIP |
| Dilution Ratio | WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 ChIP: 1/20 |
| Research Field | Epigenetics and Nuclear Signaling |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |
| Expiration Date | 12 months |

Note

Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com