

# **Anti-HMG4 Rabbit Monoclonal Antibody**

## Introduction

Multifunctional protein with various roles in different cellular compartments. May act in a redox sensitive manner. Associates with chromatin and binds DNA with a preference to non-canonical DNA structures such as single-stranded DNA. Can bent DNA and enhance DNA flexibility by looping thus providing a mechanism to promote activities on various gene promoters. Proposed to be involved in the innate immune response to nucleic acids by acting as a cytoplasmic promiscuous immunogenic DNA/RNA sensor. Negatively regulates B-cell and myeloid cell differentiation. In hematopoietic stem cells may regulate the balance between self-renewal and differentiation. Involved in negative regulation of canonical Wnt signaling.

## Product parameters

Alternative Names	HMG4; HMG-4; HMG-2a
Gene ID	3149
Ge <mark>ne N</mark> ame	HMGB3
SwissProt ID	O15347
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 23 kDa; Observed MW: 23 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-5A4F11
Form	Liquid
Concentration	See label APEXB (O
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human HMG4
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB

Dilution Ratio	WB: 1/500-1/1000
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

### Protocol

Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

### Note

1. This product is for scientific research use only.

















