

Anti-HP1 gamma Rabbit Monoclonal Antibody

Introduction

Component of heterochromatin. Recognizes and binds histone H3 tails methylated at 'Lys-9', leading to epigenetic repression. Probably involved in the repression of many genes located in euchromatin, such as E2F1, MYC and CDC25A. Involved in the formation of functional kinetochore through interaction with MIS12 complex proteins. Contributes to the conversion of local chromatin to a heterochromatin-like repressive state through H3 'Lys-9' trimethylation, mediates the recruitment of the methyltransferases SUV39H1 and/or SUV39H2 by the PER complex to the E-box elements of the circadian target genes such as PER2 itself or PER1.

Product parameters

Alternative Names	HECH; HP1-GAMMA; HP1Hs-gamma; CBX3	
Gene ID	12417	
Gene Name	Cbx3	
SwissProt ID	P23198	
Host Action Ported	Rabbit	
Reactivity	Human, Mouse, Rat	
Molecular Weight	Calculated MW: 21 kDa; Observed MW: 21 kDa	
Conjugation	Unconjugated	
Ex	-	
Em	-	
Modification	Unmodified	
Clonality	IgG	
Isotype	Monoclonal Antibody	
Clonality No.	AP-10A10F4	
Form	Liquid	
Concentration	See label	
<u>Carrier</u>	Carrier Not Free	
Immunogen	A synthetic peptide of mouse HP1 gamma	
Purification	Affinity Purified	
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.	
Application	WB, IHC-P, IP	
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20	

Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.

















