

Anti-ICOS Rabbit Monoclonal Antibody

Introduction

Enhances all basic T-cell responses to a foreign antigen, namely proliferation, secretion of lymphokines, up-regulation of molecules that mediate cell-cell interaction, and effective help for antibody secretion by B-cells. Essential both for efficient interaction between T and B-cells and for normal antibody responses to T-cell dependent antigens. Does not up-regulate the production of interleukin-2, but superinduces the synthesis of interleukin-10. Prevents the apoptosis of pre-activated T-cells. Plays a critical role in CD40-mediated class switching of immunoglobin isotypes (By similarity).

Product parameters

Alternative Names	ICOS; AILIM; CD278; Inducible T-cell costimulator; Activation-inducible lymphocyte immunomediatory molecule
Gene ID	29851
Gene Name	icos
SwissProt ID	Q9Y6W8
Host	Rabbit
Reactivity	Human
Molecular Weight	-
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-5G10D7
Form	Liquid
Concentration	See label
Carrier have Ported	Carrier Not Free
Immunogen	A synthesized peptide derived from human ICOS
Purification	Affinity Purified
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	IHC-P
Dilution Ratio	IHC: 1/100-1/200

Research Field	Immunology
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.

















