

Anti-IGF2BP2 Rabbit Monoclonal Antibody

Introduction

Binds to the 5'-UTR of the insulin-like growth factor 2 (IGF2) mRNAs. Binding is isoform-specific. May regulate translation of target mRNAs.

Product parameters

Alternative Names	IGF2BP2; IMP2; VICKZ2; Insulin-like growth factor 2 mRNA-binding protein 2; IGF2 mRNA-binding protein 2; IMP-2; Hepatocellular carcinoma autoantigen p62; IGF-II mRNA-binding protein 2; VICKZ family member 2
Gene ID	10644
Gene Name	IGF2BP2
SwissProt ID	Q9Y6M1
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 66 kDa; Observed MW: 66,62 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-4G2B11
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human IMP2
Purification	Affinity Purified
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, ICC/IF
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200
Research Field	Epigenetics and Nuclear Signaling
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C

Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

