

Anti-IL-32 Rabbit Monoclonal Antibody

Introduction

Cytokine that may play a role in innate and adaptive immune responses. It induces various cytokines such as TNFA/TNF-alpha and IL8. It activates typical cytokine signal pathways of NF-kappa-B and p38 MAPK.

Product parameters

| | |
|--------------------|---|
| Alternative Names | IL-32; IL 32alpha; IL 32beta; IL 32delta; IL 32gamma; IL32; NK4; TAlF; TAlFa; TAlFb; TAlFc; TAlFd |
| Gene ID | 9235 |
| Gene Name | IL32 |
| SwissProt ID | P24001 |
| Host | Rabbit |
| Reactivity | Human |
| Molecular Weight | Calculated MW: 27 kDa; Observed MW: 23 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-20A8A3 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Free |
| Immunogen | A synthesized peptide derived from human IL32 |
| Purification | Affinity Purified |
| Buffer System | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Application | WB |
| Dilution Ratio | WB: 1/500-1/1000 |
| Research Field | Immunology |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |

| | |
|-----------------|----------------------------------|
| Expiration Date | 12 months |
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

