

## **Anti-KDM1A (1B2) Mouse Monoclonal Antibody**

## Introduction

Histone demethylase that demethylates both 'Lys-4' (H3K4me) and 'Lys-9' (H3K9me) of histone H3, thereby acting as a coactivator or a corepressor, depending on the context. Acts by oxidizing the substrate by FAD to generate the corresponding imine that is subsequently hydrolyzed. Acts as a corepressor by mediating demethylation of H3K4me, a specific tag for epigenetic transcriptional activation. Demethylates both mono-(H3K4me1) and di-methylated (H3K4me2) H3K4me. May play a role in the repression of neuronal genes.

## Product parameters

Alternative Names	KDM1A; AOF2; KDM1; KIAA0601; LSD1; Lysine-specific histone demethylase 1A; BRAF35-HDAC complete protein BHC110; Flavin-containing amine oxidase domain-containing protein 2
Gene ID	23028
Gene Name	KDM1A
SwissProt ID	O60341
Host Achieve Perfec	Mouse
Reactivity	Human, Mouse, Monkey
Molecular Weight	Calculated MW: 93 kDa; Observed MW: 93 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	lgG1
Isotype	Monoclonal Antibody
Clonality No.	AP-13E6A8
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen Parlac	Purified recombinant fragment of human LSD1 expressed in E. Coli.
Purification	Ascitic Fluid
Buffer System	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Application	WB
Dilution Ratio	WB: 1/500-1/1000
Research Field	Epigenetics and Nuclear Signaling

Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.





