

Anti-Liver Carboxylesterase 1 Rabbit Monoclonal Antibody

Introduction

Involved in the detoxification of xenobiotics and in the activation of ester and amide prodrugs. Hydrolyzes aromatic and aliphatic esters, but has no catalytic activity toward amides or a fatty acyl-CoA ester. Hydrolyzes the methyl ester group of cocaine to form benzoylecgonine.

Product parameters

Alternative Names	ACAT; CE 1; CEH; CES1; CES2; CESDD1; Egasyn; ES-HTEL; ES-x; Es22; Esterase 22; hCE 1; HMSE; HMSE1; REH; SES1; TGH; Triacylglycerol hydrolase
Gene ID	1066
Gene Name	CES1
SwissProt ID	P23141
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 63 kDa; Observed MW: 55 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-12E4B7
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human Liver Carboxylesterase 1
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, FC, IP
Dilution Ratio	WB: 1/500-1/1000 IP: 1/40 FC: 1/50-1/100
Research Field	Cardiovascular
Product Categories	Primary antibody
Shipping	Blue ice

Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology
www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

