

## **Anti-LRRK2 Rabbit Monoclonal Antibody**

## Introduction

Positively regulates autophagy through a calcium-dependent activation of the CaMKK/AMPK signaling pathway. The process involves activation of nicotinic acid adenine dinucleotide phosphate (NAADP) receptors, increase in lysosomal pH, and calcium release from lysosomes. Together with RAB29, plays a role in the retrograde trafficking pathway for recycling proteins, such as mannose 6 phosphate receptor (M6PR), between lysosomes and the Golgi apparatus in a retromer-dependent manner.

## Product parameters

Alternative Names	AURA17; Dardarin antibody; ; Leucine rich repeat kinase 2; LRRK 2 antibody; LRRK2; LRRK2_HUMAN PARK 8; PARK8; RIPK7; ROCO 2; ROCO2
Gene ID	120892
Gene Name	LRRK2
SwissProt ID	Q5S007
Host Achieve Perfec	Rabbit
Reactivity	Human, Mouse
Molecular Weight	Calculated MW: 286 kDa; Observed MW: 286 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-17A12G8
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen Police	A synthesized peptide derived from human LRRK2
Purification	Affinity Chromatography
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, ICC/IF
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200
Research Field	Neuroscience

Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.





Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

## Note

1. This product is for scientific research use only.





