

# **Anti-MAGEC2 Rabbit Monoclonal Antibody**

### Introduction



Proposed to enhance ubiquitin ligase activity of RING-type zinc finger-containing E3 ubiquitin-protein ligases.

## Product parameters

| Alternative Names   | CT10, MAGE-C2 antigen, MAGE-E1 antigen, MAGEC2, HCA587, MAGEE1                          |
|---------------------|---|
| Gene ID             | 51438   |
| Gene Name           | MAGEC2  |
| SwissProt ID        | Q9UBF1  |
| Host                | Rabbit  |
| Reactivity          | Human, Mouse  |
| Molecular Weight    | Calculated MW: 41 kDa; Observed MW: 55 kDa  |
| Conjugation         | Unconjugated  |
| Ex /\ P             | ×BIO APE×BIO  |
| Em Achieve Perfecti | Explore the Unknown Achieve Perfection, Explore the Unknown                             |
| Modification        | Unmodified  |
| Clonality           | IgG   |
| Isotype             | Monoclonal Antibody   |
| Clonality No.       | AP-3G6C1  |
| Form                | Liquid  |
| Concentration       | See label   |
| Carrier             | Carrier Not Free  |
| Immunogen           | Recombinant protein of human MAGEE1   |
| Purification        | Affinity Purified   |
| Buffer System       | 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA. |
| Application         | WB, IHC-P, IP   |
| Dilution Ratio      | WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20   |
| Research Field      | -   |
| Product Categories  | Primary antibody  |
| Shipping            | Blue ice  |
| Storage             | -20°C   |
| Expiration Date     | 12 months   |
|                     | 1   |

#### Protocol

Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

### Note

1. This product is for scientific research use only.





