

Anti-MEK3/MEK6 Rabbit Monoclonal Antibody

Introduction

Dual specificity kinase. Is activated by cytokines and environmental stress in vivo. Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in the MAP kinase p38.

Product parameters

| | |
|--------------------|--|
| Alternative Names | MEK6; MKK6; MAPKK6; PRKMK6; SAPKK3; MAP2K6; MEK3; MAP kinase kinase 3; MAPKK3; MAPK/ERK kinase 3 |
| Gene ID | 5606/5608 |
| Gene Name | MAP2K3/MAP2K6 |
| SwissProt ID | P46734/P52564 |
| Host | Rabbit |
| Reactivity | Human, Mouse |
| Molecular Weight | Calculated MW: 39,37 kDa; Observed MW: 39,37 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-4H9B3 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Not Free |
| Immunogen | A synthesized peptide derived from human MEK3/MEK6 |
| Purification | Affinity Purified |
| Buffer System | 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA. |
| Application | WB, ICC/IF, IP |
| Dilution Ratio | WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/20 |
| Research Field | Signal Transduction |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |

| | |
|-----------------|----------------------------------|
| Expiration Date | 12 months |
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APEx BIO Technology
www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

