

Anti-MELK Rabbit Monoclonal Antibody

Introduction

Phosphorylates ZNF622 and may contribute to its redirection to the nucleus. May be involved in the inhibition of spliceosome assembly during mitosis.

Product parameters

| | |
|--------------------|--|
| Alternative Names | MELK; KIAA0175; Maternal embryonic leucine zipper kinase; hMELK; Protein kinase Eg3; pEg3 kinase; Protein kinase PK38; hPK38; Tyrosine-protein kinase MELK |
| Gene ID | 9833 |
| Gene Name | MELK |
| SwissProt ID | Q14680 |
| Host | Rabbit |
| Reactivity | Human, Rat |
| Molecular Weight | Calculated MW: 75 kDa; Observed MW: 75 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-17G4F9 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Not Free |
| Immunogen | A synthetic peptide of human MELK |
| Purification | Affinity Purified |
| Buffer System | 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA. |
| Application | WB |
| Dilution Ratio | WB: 1/500-1/1000 |
| Research Field | Signal Transduction |
| Product Categories | Primary antibody |
| Shipping | Blue ice |
| Storage | -20°C |

| | |
|-----------------|----------------------------------|
| Expiration Date | 12 months |
| Note | Please avoid freeze-thaw cycles. |

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APEx BIO Technology
www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

