

Anti-MERTK Rabbit Monoclonal Antibody

Introduction

MERTK signaling plays a role in various processes such as macrophage clearance of apoptotic cells, platelet aggregation, cytoskeleton reorganization and engulfment. Functions in the retinal pigment epithelium (RPE) as a regulator of rod outer segments fragments phagocytosis. Plays also an important role in inhibition of Toll-like receptors (TLRs)-mediated innate immune response by activating STAT1, which selectively induces production of suppressors of cytokine signaling SOCS1 and SOCS3.

Product parameters

Alternative Names	MERTK; MER; Tyrosine-protein kinase Mer; Proto-oncogene c-Mer; Receptor tyrosine kinase MerTK
Gene ID	10461
Gene Name	MERTK
SwissProt ID	Q12866
Host	Rabbit
Reactivity	Human
Molecular Weight	Calculated MW: 110 kDa; Observed MW: 210 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-12G7H4
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human MERTK
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB, IHC-P, IP
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IP: 1/20
Research Field	Cell Biology

Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.



Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

