

Anti-MLLT6 Rabbit Monoclonal Antibody

Introduction



Enables histone binding activity and nucleosome binding activity. Predicted to be involved in positive regulation of transcription by RNA polymerase II. Predicted to act upstream of or within several processes, including negative regulation of histone H3-K79 methylation; renal potassium excretion; and renal sodium excretion. Predicted to be active in nucleus.

Product parameters

Alternative Names	AF17
Gene ID	4302
Gene Name	MLLT6
SwissProt ID	P55198
Host	Rabbit
Reactivity	Human () A PEXBIO
Molecular Weight	Calculated MW: 112 kDa; Observed MW: 112 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-10G9D5
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Recombinant protein of human MLLT6
Pu <mark>rifica</mark> tion	Affinity Purified Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB
Dilution Ratio	WB: 1/500-1/1000
Research Field	-
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol



Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





