

Anti-MMP12 Rabbit Monoclonal Antibody

Introduction

MMP-12 lacks a transmembrane domain and furin cleavage site. The zymogen for MMP-12 is about 54 kDa, and is quickly activated to the 45 kDa form, and this breaks down to cascade of active forms, ending with the common 22 kDa form. Stimulated macrophages produce MMP-12; it has also been found in osteosarcoma cells, synovial fibroblasts and lung fibroblasts.

Product parameters

Alternative Names	Macrophage metalloelastase; MME; 3.4.24.65; Macrophage elastase; ME; hME; Matrix metalloproteinase-12; MMP-12; MMP12; HME
Gene ID	4321
Gene Name	MMP12
SwissProt ID	P39900
Host	Rabbit
Reactivity	Human, Rat, Hamster
Molecular Weight	Calculated MW: 54 kDa; Observed MW: 54,45 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-16C6H11
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	A synthetic peptide of human MMP12
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB, IP
Dilution Ratio	WB: 1/500-1/1000 IP: 1/20
Research Field	Cardiovascular
Product Categories	Primary antibody

Shipping	Blue ice
Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol



Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology
www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

