

Anti-MTHFR Rabbit Monoclonal Antibody

Introduction



The protein encoded by this gene catalyzes the conversion of 5,10-methylenetetrahydrofolate to 5-methyltetrahydrofolate, a co-substrate for homocysteine remethylation to methionine. Genetic variation in this gene influences susceptibility to occlusive vascular disease, neural tube defects, colon cancer and acute leukemia, and mutations in this gene are associated with methylenetetrahydrofolate reductase deficiency.

Product parameters

| Alternative Names | MTHFR; Methylenetetrahydrofolate reductase |
|--------------------|---|
| Gene ID | 4524 |
| Gene Name | MTHFR |
| SwissProt ID | P42898 |
| Host | Rabbit |
| Reactivity | Human, Mouse |
| Molecular Weight | Calculated MW: 75 kDa; Observed MW: 75 kDa |
| Conjugation | Unconjugated |
| Ex | - |
| Em | - |
| Modification | Unmodified |
| Clonality | IgG |
| Isotype | Monoclonal Antibody |
| Clonality No. | AP-4C12D8 |
| Form | Liquid |
| Concentration | See label |
| Carrier | Carrier Not Free |
| Immunogen | Recombinant protein of human MTHFR |
| Purification | Affinity Purified Affinity Purified |
| Buffer System | 50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA. |
| Application | WB, ICC/IF, IP |
| Dilution Ratio | WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/20 |
| Research Field | Signal Transduction |
| Product Categories | Primary antibody |

| Shipping | Blue ice |
|-----------------|----------------------------------|
| Storage | -20°C |
| Expiration Date | 12 months |
| Note | Please avoid freeze-thaw cycles. |

| Protocol



Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





