

Anti-NAT10 Rabbit Monoclonal Antibody

Introduction



Has protein acetyltransferase activity in vitro. Can acetylate both histones and microtubules. Histone acetylation may regulate transcription and mitotic chromosome de-condensation.

Product parameters

Alternative Names	ALP; hALP; N acetyltransferase like protein; NAT10; NET43
Gene ID	55226
Gene Name	NAT10
SwissProt ID	Q9H0A0
Host	Rabbit
Reactivity	Human, Rat
Molecular Weight	Calculated MW: 116 kDa; Observed MW: 116 kDa
Conjugation	Unconjugated A PEXBLO
Ex Achieve Perfect	Explore the Unknown Actieve Perfection, Explore the Unknown
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-14F1F10
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Recombinant protein of human NAT10
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Ap <mark>plica</mark> tion	WB, ICC/IF, IP
Dilution Ratio	WB: 1/500-1/1000 IF: 1/50-1/200 IP: 1/20
Research Field	Cell Biology
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C
	.1

Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





