

Anti-NCF1 Rabbit Monoclonal Antibody



NCF2, NCF1, and a membrane bound cytochrome b558 are required for activation of the latent NADPH oxidase (necessary for superoxide production).

Product parameters

Alternative Names	NCF1; NCF1A; NOXO2; SH3PXD1A; p47phox
Gene ID	653361
Gene Name	NCF1
SwissProt ID	P14598
Host	Rabbit
Reactivity	Human
Molecular Weight	Calculated MW: 45 kDa; Observed MW: 45 kDa
Conjugation	Unconjugated PEX BIO
Ex Actueve Perfect	Explore the Unknown
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-14B4B2
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human NCF1
Purification	Affinity Purified
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol
Application	
Dilution Ratio	WB: 1/500-1/1000 IP: 1/50
Research Field	-
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C

Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.





