

Anti-Niemann Pick C2 Rabbit Monoclonal Antibody

Introduction

This gene encodes a protein containing a lipid recognition domain. The encoded protein may function in regulating the transport of cholesterol through the late endosomal/lysosomal system. Mutations in this gene have been associated with Niemann-Pick disease, type C2 and frontal lobe atrophy. [provided by RefSeq, Jul 2008]

Product parameters

Alternative Names	HE1; EDDM1
Gene ID	10577
Gene Name	NPC2
SwissProt ID	P61916
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 17 kDa; Observed MW: 17 kDa
Conjugation	Unconjugated
Ex	-
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-10C11E1
Form	Liquid
Concentration	See label
Carrier	Carrier Not Free
Immunogen	Recombinant protein of human Niemann Pick C2
Purification	Affinity Purified
Buffer System	50mM Tris-Glycine (pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% BSA.
Application	WB, IHC-P
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100
Research Field	Cardiovascular
Product Categories	Primary antibody
Shipping	Blue ice

Storage	-20°C
Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

Protocol

Configure the product according to the application range and recommended dilution ratio.

***Note:** The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

Note

1. This product is for scientific research use only.



APExBIO Technology

www.apexbt.com

7505 Fannin street, Suite 410, Houston, TX 77054.

Tel: +1-832-696-8203 | Fax: +1-832-641-3177 | Email: info@apexbt.com

