

# **Anti-NUDC Rabbit Monoclonal Antibody**

## Introduction

cessary for correct formation of mitotic

Plays a role in neurogenesis and neuronal migration (By similarity). Necessary for correct formation of mitotic spindles and chromosome separation during mitosis. Necessary for cytokinesis and cell proliferation.

## Product parameters

Alternative Names	HNUDC; MNUDC; NPD011; nudC; SIG92
Gene ID	10726
Gene Name	NUDC
SwissProt ID	Q9Y266
Host	Rabbit
Reactivity	Human, Mouse, Rat
Molecular Weight	Calculated MW: 38 kDa; Observed MW: 42 kDa
Conjugation	Unconjugated APEXE
Ex Achieve Perfecti	4), Explore the Unknown.  Achieve Perfection, Explore the Unknown.
Em	-
Modification	Unmodified
Clonality	IgG
Isotype	Monoclonal Antibody
Clonality No.	AP-15E10C1
Form	Liquid
Concentration	See label
Carrier	Carrier Free
Immunogen	A synthesized peptide derived from human NUDC
Purification	Affinity Purified
Buffer System	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Application	WB, IHC-P, ICC/IF, FC, IP
Dilution Ratio	WB: 1/500-1/1000 IHC: 1/50-1/100 IF: 1/50-1/200 IP: 1/50 FC: 1/50-1/100
Research Field	-
Product Categories	Primary antibody
Shipping	Blue ice
Storage	-20°C

Expiration Date	12 months
Note	Please avoid freeze-thaw cycles.

#### Protocol

Configure the product according to the application range and recommended dilution ratio.

\*Note: The primary antibody dilution buffer options: WB - Primary Antibody Dilution Buffer (Cat. #: K1200, Not for HRP/AP conjugated antibodies), Immunostaining - Immunol Staining Primary Antibody Dilution Solution (Cat. #: K4655).

#### Note

1. This product is for scientific research use only.





